

A possible route to spontaneous reduction of the heat conductivity by a temperature gradient driven instability in electron-ion plasmas

Makoto Hattori

Astronomical Institute, Tôhoku University, Aoba, Sendai 980-8578, Japan; hattori@astr.tohoku.ac.jp

and

Keiichi Umetsu

Astronomical Institute, Tôhoku University, Aoba, Sendai 980-8578, Japan; keiichi@astr.tohoku.ac.jp

ABSTRACT

We have shown that there exists low-frequency growing modes driven by a global temperature gradient in electron and ion plasmas, by linear perturbation analysis within the frame work of plasma Kinetic theory. The driving force of the instability is the local deviation of the distribution function from the Maxwell-Boltzmann due to global temperature gradient. Application to the intracluster medium shows that scattering of the particles due to waves excited by the instability is possible to reduce mean free paths of electron and ion down to five to seven order of magnitude than the mean free paths due to Coulomb collisions. This may provide a hint to explain why hot and cool gas can co-exist in the intracluster medium in spite of the very short evaporation time scale due to thermal conduction if the conductivity is the classical Spitzer value. Our results suggest that the realization of the global thermal equilibrium is postponed by the local instability which is induced for quicker realization of local thermal equilibrium state in plasmas. The instability provides a new possibility to create and grow cosmic magnetic fields without any seed magnetic field.

Subject headings: cooling flows — instabilities — plasmas — galaxies: clusters: general

1. Introduction

One of the most important results obtained by Japanese/US X-ray satellite ASCA (Tanaka, Inoue, & Holt 1994) is the first spectro-scopical confirmation of existence of at least two different temperature X-ray emitting hot gas in the cluster central region (Fukazawa et al. 1994; Ikebe 1995; Ikebe et al. 1999); the cooler is about 1keV and the hotter is about 2keV–10keV. Although the interpretation of the result is still controversial (Fabian 1994; Ikebe et al. 1999), there is a common feature among the different interpretations, that is, a cool region with temperature T_1 is surrounded by a hot region with temperature T_2 maintaining a pressure equilibrium (see §6.2). Since the total thermal energy in the hot region is much higher than that in the cool region, the hot region can be treated as a heat reservoir. Suppose that the interface separating two plasma regions with different temperatures has a thickness of L and temperature variation in the interface region is almost linear in spatial scale. Assume that the electron mean free path λ_e is smaller than L . The time scale τ_{ev} of the evaporation due to heat conduction can be roughly estimated by a time scale for electrons in the hotter region to diffuse across the interface, and is given as

$$\tau_{ev} \sim (\lambda_e/v_{th,e}) \times (L/\lambda_e)^2 = \epsilon^{-1}(L/v_{th,e}) \sim 10^5(L/4\text{kpc})(k_B T_e/4\text{keV})^{-0.5}(\epsilon/1.0)^{-1}\text{yr}, \quad (1)$$

where $v_{\text{th},e} \equiv \sqrt{2k_B T_e / m_e}$ is the electron thermal velocity, T_e is the electron temperature, and m_e is the electron mass; ϵ is defined by $\epsilon \equiv \lambda_e / L$. In the cluster central region, the electron mean free path due to Coulomb collision is obtained as $\lambda_C \sim 0.5(k_B T_e / 4\text{keV})^2 (n_e / 10^{-2}\text{cm}^{-3})^{-1} \text{kpc}$ (Sarazin 1988). Therefore, the time scale of the evaporation is about 10^6 yr and is much shorter than the age of cluster of galaxies, which is about the Hubble time. For maintaining the interface at least as long as the age of the cluster, the reduction of the electron mean free path in four order of magnitude from the Coulomb mean free path to realize ϵ of $\sim 10^{-5}$ is required. Soker & Sarazin (1990) and Makishima (1997) proposed that the cool and hot phases are thermally insulated from each other by magnetic fields. However, their models involve another problem how such special magnetic field structures are obtained in the cluster central region. In this paper, we show that there is an alternative possibility to reduce electron mean free path in many order of magnitude without any special assumption on magnetic field structure.

Ramani & Laval (1978) found a new plasma instability which may relate to the reduction of the electron mean free path, henceforce the reduction of the heat conduction. They showed that the temperature gradient leads to an anisotropic electron velocity distribution function and that the anisotropy of the velocity distribution function drives the instability like the Weibel instability (Weibel 1959; Fried 1959; Melrose 1986). They proposed that the chaotic magnetic field and electric field, produced by the plasma wave induced by the instability, scatter electrons, so that the electron mean free path is reduced in many order of magnitude. However, the instability found by them cannot be applied to reduce the electron mean free path in many order of magnitude because of the following reasons. Since they assumed that only electrons respond to the mode and ions were treated as fixed back ground particles, the phase velocity of the wave must be faster than the ion sound velocity $v_{\text{th},i} = \sqrt{2k_B T_i / m_i}$, where T_i , n_i , and m_i are the ion temperature, ion number density, and ion mass, respectively. In the astrophysical plasma, the ion mass can be safely replaced by the proton mass. The unstable mode found by Ramani & Laval (1978) has non zero real part of the wave frequency with a phase velocity of $\sim \epsilon v_{\text{th},e}$. Therefore, the application limit of their analysis sets a relatively high lower limit on ϵ as $\epsilon > \sqrt{m_e / m_i} \sim 0.025$ when $T_i = T_e$ (hereafter we refer to this lower limit as the wall of the square root of the mass ratio). To explain the two-phase nature of the hot gas in cluster central region, the instability found by them is practically not useful and the mechanism which can break the wall of the square root of the mass ratio is required.

The first application of the Ramani & Laval (1978) type instability to the astrophysical plasma was made by Levinson & Eichler (1992). They extend the Ramani & Laval (1978)'s analysis to include the non-zero background magnetic field and found the low-frequency unstable mode, similar to the Ramani & Laval (1978) type modes. Pistnner, Levinson & Eichler (1996) first applied the Levinson & Eichler (1992)'s results to the cluster cooling flow. However, they applied their results to reduce electron mean free path down to 10^{-3} and far less than the lower limit, bounded by the application limit of the wall of the square root of the mass ratio. Further they neglected the cluster gravitational force in their Boltzmann equation in spite of taking into account a pressure gradient with a scale height comparable to the cluster gravitational scale height. As a result, they implicitly assumed the existence of the steady electric field for the electric force acting on an electron to balance with the pressure gradient force acting on the electron. Since the pressure gradient force acting on an ion balances with the cluster gravitational force acting on the ion, the electric force acting on an ion is as strong as the cluster gravitational force acting on the ion. The existence of such a strong electric field in cluster of galaxies is unlikely. Further, the model adopted as the temperature distribution in the cluster central region obtained by White & Sarazin (1987) only describes the temperature distribution of the hottest phase gas. Therefore, the model adopted by them is inappropriate to answer why the multi-phase gas clumps can co-exist in the cluster of galaxies.

In this paper, we extend the Ramani & Laval (1978)’s analysis including the response of the ion to examine whether we can break the wall of the square root of the mass ratio. We limit our attention to the pressure equilibrium plasma; in other words, the scale of the interface is much smaller than the pressure scale height. Then, the gravitational force can be neglected. The plan of the paper is as follows. The readers who are not interested in the details of the derivation of the dispersion relation can skip §§3 and 4. In §2, we set the model, and summarize the basic assumptions and the application limit coming from the assumptions. In §3, we deduce the solution of the Boltzmann equation under the situations set in §2 using the Chapman-Enskog expansion of the electron and ion velocity distribution functions. We deduce the dispersion relations under low-frequency conditions in §4. In §5, we summarize the characteristics of the instability and modes found in §4. The application of the results to the hot plasma in the central region of cluster of galaxies is discussed in §6. Finally, §7 summarizes another important implications from our results and remained problems.

2. The model and application limit

Consider the situation described in Fig. 1. Two plasma regions with different temperatures contact maintaining pressure equilibrium. The electron temperature in the transition region with a depth of L varies almost linearly from T_1 to T_2 ($T_1 < T_2$) along the x -direction: $\nabla T(x)/T(x) = \delta_T L^{-1}$ where $\delta_T \equiv (T_2 - T_1)/T$. In clusters of galaxies, the pressure scale height is about the core radius of the cluster mass distribution, say $\sim 50\text{kpc}$ (e.g., Hattori, Kneib, & Makino 1999). Therefore, if the size of the interface is smaller than $\sim 10\text{kpc}$, the pressure equilibrium condition for the interface region can be safely applied. The ion temperature is assumed to be the same as electron temperature, $T_i = T_e$. The mean free paths λ_e and λ_i of electron and ion, respectively, are assumed to be defined by the scattering due to plasma waves. If the scattering of the particle due to electric field E of the wave is the dominant source, the mean free path of each particle could be described as follows. Assume that the electric field associated with plasma wave is coherent in the scale of order of the wave length λ and is chaotic over the scale. Then the momentum change of the particles can be described by the random walk manner in the momentum space. The mean free path is defined as the length scale that the particles gain or loss its momentum comparable to their original momentum during they move the scale. The momentum change of electron during one step is $m_e \Delta v_e \sim eE\lambda/v_{\text{th},e}$. The number of steps required for electron to gain or loss their momentum comparable to its original momentum is $[m_e v_{\text{th},e}/(eE\lambda/v_{\text{th},e})]^2 \sim (2k_B T_e/eE\lambda)^2$. Then we obtain

$$\lambda_e \sim \left(\frac{2k_B T_e}{eE\lambda} \right)^2 \lambda \sim \left(\frac{2k_B T_e}{eE} \right)^2 \frac{1}{\lambda}. \quad (2)$$

Similarly, the mean free path for ion can be obtained as

$$\lambda_i \sim \left(\frac{2k_B T_i}{eE\lambda} \right)^2 \lambda \sim \left(\frac{2k_B T_i}{eE} \right)^2 \frac{1}{\lambda}. \quad (3)$$

Since we are now assuming that electron and ion have the same temperature, the electron and ion mean free paths due to the scattering by the wave electric field are identical. When the chaotic magnetic field associated with the plasma wave is the dominant source of the particle scattering, the particle mean free path would be defined in the following way. The magnetic moment of the particles, $\mu = (1/2m_e v_\perp^2)/B$, is a conserved variable in a magnetized plasma, where $v_\perp$ is the particle velocity perpendicular to the magnetic field, written as $m_e v_\perp^2 \sim 2k_B T_e$, just after the perturbed magnetic field is induced. As the magnetic-field strength gets stronger, $v_\perp$ increases to conserve μ . The maximum allowable value of $v_\perp$ is $\sqrt{3k_B T_e/m_e}$,

since the kinetic energy of the particle must be conserved. Then it follows that an electron at the place with a magnetic field strength of B_1 is scattered back at the place with a magnetic field strength of $B_{\text{cr}} \sim 3/2 B_1$ owing to the mirror effect. The mean free path is then defined as the length scale along the magnetic field from the place where the magnetic field strength is B_1 to the place where the magnetic field strength gets B_{cr} . Since the critical value B_{cr} of the magnetic field strength, in other words the place of the bouncing, does not depend on the particle species and only depends on their initial position and the magnetic-field configuration. Therefore, the mean free paths of ion and electron must be also the same in this case. These arguments show that the mean free paths of electron and ion due to scattering by plasma waves may have the same order of magnitude. Therefore, we assume that electrons and ions have a similar value of ϵ .

In this paper, perturbations with a short-wavelength wave, such that

$$\lambda \ll \lambda_e, \quad (4)$$

are treated. Therefore, the collision term in the Boltzmann equation for the perturbed variables can be neglected.

Self-consistency check of the obtained results with all these conditions constrain the allowable range of ϵ . In addition, two further conditions provide important limits on ϵ . First of all, the electron mean free path due to Coulomb collision must be shorter than L at least before the onset of the instability, otherwise there is no chance for the electron distribution function to have anisotropy as described in the next section, and hence no chance to have instability. This condition yields an lower limit on L as

$$L > 0.5(k_B T_e / 4\text{keV})^2 (n_e / 10^{-2} \text{cm}^{-3})^{-1} \text{kpc}. \quad (5)$$

Secondly, the growth time scale of the unstable mode, ω_i^{-1} , must be shorter than the time scale of thermal diffusion when waves do not exist, in order for the waves to grow before the temperature difference will be erased owing to quick thermal conduction. This condition provides

$$\omega_i^{-1} < \epsilon^{-1} L / v_{\text{th},e}, \quad (6)$$

where only the case that the electron mean free path is shorter than L is considered. A further condition has to be required when applied to the clusters, that is the growth time scale must be shorter than the age of the object, conservatively the age of the universe:

$$\omega_i^{-1} < 1/H_0 = 10^{10} h^{-1} \text{yr}, \quad (7)$$

where $H_0 = 100 h \text{km/sec/Mpc}$ is the Hubble constant.

3. Distribution Function

When $\epsilon < 1$, the discussion given in the previous section ensures that the solution of the Boltzmann equation of electron and ion can be expanded in powers of ϵ (Chapman & Cowling 1960)

$$f_e = f_{\text{m},e} + f_e^{(1)} + f_e^{(2)} + \dots, \quad (8)$$

$$f_i = f_{\text{m},i} + f_i^{(1)} + f_i^{(2)} + \dots, \quad (9)$$

where $f_{\text{m},e}$ and $f_{\text{m},i}$ are Maxwellian distribution functions for electron and ion, respectively; $f_e^{(k)}$ and $f_i^{(k)}$ ($k = 1, 2, \dots$) describe the deviation of distribution functions from the Maxwellian in

order of ϵ^k for electron and ion, respectively. Since we assume that $T_i(x) = T_e(x) \equiv T(x)$ and the charge neutrality must be maintained, the Maxwellian parts for both electron and ion are written as $f_{m,e} = n_0(x)[\pi v_{th,e}(x)]^{-3/2} \exp\{-[v/v_{th,e}(x)]^2\}$ and $f_{m,i} = n_0(x)[\pi v_{th,i}(x)]^{-3/2} \exp\{-[v/v_{th,i}(x)]^2\}$, respectively. Here $n_0(x)$ is the electron number density, $v_{th,e}(x) \equiv \sqrt{2k_B T(x)/m_e}$, and $v_{th,i}(x) \equiv \sqrt{2k_B T(x)/m_i}$.

The pressure equilibrium assumption gives

$$\frac{\nabla n_0}{n_0} = -\frac{\nabla T}{T} = -\frac{1}{L}\delta T. \quad (10)$$

Once this condition is satisfied, the pressure is time independent everywhere, even if a secular variation of the temperature due to thermal conduction is taken into account. Then the time dependence of the density can be related to that of the temperature as

$$\frac{1}{n_0} \frac{\partial n_0}{\partial t} = -\frac{1}{T} \frac{\partial T}{\partial t}. \quad (11)$$

The Boltzmann equation under the pressure equilibrium condition without background electric and magnetic fields leads to

$$\frac{\partial f_e}{\partial t} + v_x \frac{\partial f_e}{\partial x} = \left(\frac{\partial f_e}{\partial t} \right)_c, \quad (12)$$

where $(\partial f/\partial t)_c$ is the collision term and the equilibrium electric field does not appear because of the pressure equilibrium condition. The secular time variation of the distribution function is ascribed to the secular variation of the temperature due to thermal conduction. Since the cool region with a temperature of T_1 is considered to be immersed in a heat bath with a temperature of T_2 , the temperature of the interface increases monotonically owing to the heat conduction. Hence, the time evolution of temperature can be described as

$$\frac{\partial T}{\partial t} = \epsilon \delta_T \frac{v_{th,e}}{L} T. \quad (13)$$

Then,

$$\left| \frac{\partial f_e}{\partial t} \right| \sim \frac{1}{T} \left| \frac{\partial T}{\partial t} \right| f_e \sim \epsilon \delta_T \frac{v_{th,e}}{L} f_e. \quad (14)$$

The simplest choice of the collision term is the Krook operator, such that

$$\left(\frac{\partial f_e}{\partial t} \right)_c = -\nu_e (f_e - f_{m,e}), \quad (15)$$

where $\nu_e = v_{th,e}/\lambda_e$ is the collision frequency assumed to be constant. Substituting the collision term by the Krook operator, ordering the Boltzmann equation in ϵ^1 and ϵ^2 provides

$$f_e^{(1)} = -\frac{1}{\nu_e} v_x \frac{\partial f_{m,e}}{\partial x}, \quad (16)$$

$$f_e^{(2)} = -\frac{1}{\nu_e} \left(\frac{\partial f_{m,e}}{\partial t} + v_x \frac{\partial f_e^{(1)}}{\partial x} \right). \quad (17)$$

It shows that the system can be treated as steady state in the first order of ϵ but the secular variation cannot be negligible in the second order of ϵ . The time derivative of the Maxwell-Boltzmann distribution function can be described as

$$\frac{\partial f_{m,e}}{\partial t} = -\frac{1}{2} \left(5 - \frac{2v^2}{v_{th,e}^2} \right) \frac{1}{T} \frac{\partial T}{\partial t} f_{m,e} = -\frac{\epsilon \delta_T}{2} \frac{v_{th,e}}{L} \left(5 - \frac{2v^2}{v_{th,e}^2} \right) f_{m,e}. \quad (18)$$

The first- and the second-order solutions for the electron distribution function are thus obtained as follows:

$$f_e^{(1)} = -\epsilon\delta_T \frac{v_x}{v_{\text{th},e}} \left(5 - \frac{m_e v^2}{k_B T}\right) f_{\text{m},e}, \quad (19)$$

$$f_e^{(2)} = \frac{\epsilon^2 \delta_T}{2} \left[\left(5 - \frac{2v^2}{v_{\text{th},e}^2}\right) + \frac{v_x^2}{v_{\text{th},e}^2} \left(5 - \frac{4v^2}{v_{\text{th},e}^2}\right) + \frac{v_x^2}{2v_{\text{th},e}^2} \left(5 - \frac{2v^2}{v_{\text{th},e}^2}\right)^2 \right] f_{\text{m},e}, \quad (20)$$

where we have neglected the second derivative of $T(x)$ with respect to x in derivation of the second-order solution.

These equations show that the anisotropy in velocity distribution functions are induced by the existence of the temperature gradient. The first-order solution is an odd function of the velocity. On the other hand, the second-order solution is even in velocity. It is straight forward to make sure that the zero background electric field assumption is consistent with the zero electric current condition under the pressure equilibrium plasma condition, by checking $\langle f_e \vec{v} \rangle = \vec{0}$, where $\langle \rangle$ denotes the average over the velocity.

By substituting m_e to m_i and ν_e to $\nu_i = \nu_i/\lambda_i$ in equations (19) and (20), the first and the second order distribution functions $f_i^{(1)}$ and $f_i^{(2)}$ for ion are obtained.

4. Dispersion relation

In the short-wavelength limit, the collisionless Boltzmann equation is a good representative for the evolution of the perturbed distribution functions, and the plain wave $\exp[i(\vec{k} \cdot \vec{r} - \omega t)]$ is a good representative for the perturbed quantities (e.g., electron distribution function as $\delta f_e = f_{\vec{k},e} \exp[i(\vec{k} \cdot \vec{r} - \omega t)]$, electric field as $\vec{E} = \vec{E}_{\vec{k}} \exp[i(\vec{k} \cdot \vec{r} - \omega t)]$, and magnetic field as $\vec{B} = \vec{B}_{\vec{k}} \exp[i(\vec{k} \cdot \vec{r} - \omega t)]$). From the linearized Boltzmann equation in terms of the perturbed quantities of both electron and ion, the equations for the perturbed distribution functions of electron and ion are obtained as

$$-i(\omega - \vec{k} \cdot \vec{v}) f_{\vec{k},e} = +\frac{e}{m_e} \left(\vec{E}_{\vec{k}} + \vec{v} \times \frac{\vec{B}_{\vec{k}}}{c} \right) \cdot \left(\frac{\partial f_e}{\partial \vec{v}} \right) \quad (21)$$

$$-i(\omega - \vec{k} \cdot \vec{v}) f_{\vec{k},i} = -\frac{e}{m_i} \left(\vec{E}_{\vec{k}} + \vec{v} \times \frac{\vec{B}_{\vec{k}}}{c} \right) \cdot \left(\frac{\partial f_i}{\partial \vec{v}} \right). \quad (22)$$

As shown below, the low-frequency condition requires that the charge density be small, implying that the electric current has to be almost transverse for purely electronic oscillations. Therefore, only two modes are expected. Without loss of generality, the wave vector $\vec{k}$ can be taken in the x - y plane as $k_x = k \cos \theta$, $k_y = k \sin \theta$, and $k_z = 0$. We expect one mode with the magnetic field along the z -direction (mode (1): $B_x = B_y = 0$ and $B_z \neq 0$), and the other one with the magnetic field in the x - y plane (mode (2): $B_x, B_y \neq 0$ and $B_z = 0$). We thus denote the component of a vector along the direction of the wave vector by a k -subscript and the component along the perpendicular direction of the wave vector in the x - y plain by a $\perp$ -subscript. The x -component of velocity, for example, can be expressed as $v_x = v_k \cos \theta - v_{\perp} \sin \theta$ in terms of v_k and $v_{\perp}$.

The Faraday's law in Maxwell's equations leads to $E_z = 0$ (i.e., $\vec{E}_{\vec{k}} = (E_k, E_{\perp}, 0)$) and $kE_{\perp} = (\omega/c)B_z$ for mode (1). For mode (2), $\text{div} \vec{B} = 0$ yields $B_k = 0$ (i.e., $\vec{B}_{\vec{k}} = (0, B_{\perp}, 0)$), and the Faraday's law and

$B_z = 0$ lead to $E_\perp = 0$ and $kE_z = -(\omega/c)B_\perp$. Then we obtain

$$(\vec{k} \cdot \vec{v} - \omega)f_{\vec{k},e} = -\frac{ie}{m_e} \left[\left(E_k + \frac{v_\perp}{c} B_z \right) \frac{\partial f_e}{\partial v_k} + \left(E_\perp - \frac{v_k}{c} B_z \right) \frac{\partial f_e}{\partial v_\perp} \right], \quad (23)$$

$$(\vec{k} \cdot \vec{v} - \omega)f_{\vec{k},i} = +\frac{ie}{m_i} \left[\left(E_k + \frac{v_\perp}{c} B_z \right) \frac{\partial f_i}{\partial v_k} + \left(E_\perp - \frac{v_k}{c} B_z \right) \frac{\partial f_i}{\partial v_\perp} \right] \quad (24)$$

for mode (1), and

$$(\vec{k} \cdot \vec{v} - \omega)f_{\vec{k},e} = -\frac{ie}{m_e} \left[\left(E_k - \frac{v_z}{c} B_\perp \right) \frac{\partial f_e}{\partial v_k} + \left(E_z + \frac{v_k}{c} B_\perp \right) \frac{\partial f_e}{\partial v_z} \right], \quad (25)$$

$$(\vec{k} \cdot \vec{v} - \omega)f_{\vec{k},i} = +\frac{ie}{m_i} \left[\left(E_k - \frac{v_z}{c} B_\perp \right) \frac{\partial f_i}{\partial v_k} + \left(E_z + \frac{v_k}{c} B_\perp \right) \frac{\partial f_i}{\partial v_z} \right] \quad (26)$$

for mode (2).

To first order in $\omega/(kv_{\text{th},e})$ and $\omega/(kv_{\text{th},i})$, $(\vec{k} \cdot \vec{v} - \omega)^{-1} = (1/k)P(1/v_k) + (\omega/k^2)P(1/v_k^2) + i(\pi/k)\delta(v_k) - i(\pi/k)(\omega/k)(d\delta(v_k)/dv_k)$, where P denotes the principal value and the signs in front of the delta functions reflect the causality condition. Then, with the help of the Faraday's law, equations (23) and (24) yield

$$\begin{aligned} f_{\vec{k},e} = & -\frac{ie}{m_e} \left\{ \left[\frac{1}{k}P\frac{1}{v_k} + \frac{\omega}{k^2}P\frac{1}{v_k^2} + \frac{i\pi}{k}\delta(v_k) - \frac{i\pi\omega}{k^2}\frac{d\delta(v_k)}{dv_k} \right] \times \right. \\ & \left(E_k + \frac{v_\perp}{c} B_z \right) \left(-\frac{2v_k}{v_{\text{th},e}^2} + \frac{\epsilon\delta_T}{2v_{\text{th},e}} \left[\cos\theta \left(5 - \frac{2v^2}{v_{\text{th},e}^2} \right) - \frac{2v_k}{v_{\text{th},e}^2} (v_k \cos\theta - v_\perp \sin\theta) \left(7 - \frac{2v^2}{v_{\text{th},e}^2} \right) \right] \right) \\ & \left. - \frac{B_z}{ck} \left(-\frac{2v_\perp}{v_{\text{th},e}^2} + \frac{\epsilon\delta_T}{2v_{\text{th},e}} \left[-\sin\theta \left(5 - \frac{2v^2}{v_{\text{th},e}^2} \right) - \frac{2v_\perp}{v_{\text{th},e}^2} (v_k \cos\theta - v_\perp \sin\theta) \left(7 - \frac{2v^2}{v_{\text{th},e}^2} \right) \right] \right) \right\} f_{\text{m},e}, \end{aligned} \quad (27)$$

$$\begin{aligned} f_{\vec{k},i} = & +\frac{ie}{m_i} \left\{ \left[\frac{1}{k}P\frac{1}{v_k} + \frac{\omega}{k^2}P\frac{1}{v_k^2} + \frac{i\pi}{k}\delta(v_k) - \frac{i\pi\omega}{k^2}\frac{d\delta(v_k)}{dv_k} \right] \times \right. \\ & \left(E_k + \frac{v_\perp}{c} B_z \right) \left(-\frac{2v_k}{v_{\text{th},i}^2} + \frac{\epsilon\delta_T}{2v_{\text{th},i}} \left[\cos\theta \left(5 - \frac{2v^2}{v_{\text{th},i}^2} \right) - \frac{2v_k}{v_{\text{th},i}^2} (v_k \cos\theta - v_\perp \sin\theta) \left(7 - \frac{2v^2}{v_{\text{th},i}^2} \right) \right] \right) \\ & \left. - \frac{B_z}{ck} \left(-\frac{2v_\perp}{v_{\text{th},i}^2} + \frac{\epsilon\delta_T}{2v_{\text{th},i}} \left[-\sin\theta \left(5 - \frac{2v^2}{v_{\text{th},i}^2} \right) - \frac{2v_\perp}{v_{\text{th},i}^2} (v_k \cos\theta - v_\perp \sin\theta) \left(7 - \frac{2v^2}{v_{\text{th},i}^2} \right) \right] \right) \right\} f_{\text{m},i} \end{aligned} \quad (28)$$

for mode (1). Similarly, from equations (25) and (26), we obtain

$$\begin{aligned} f_{\vec{k},e} = & -\frac{ie}{m_e} \left\{ \left[\frac{1}{k}P\frac{1}{v_k} + \frac{\omega}{k^2}P\frac{1}{v_k^2} + \frac{i\pi}{k}\delta(v_k) - \frac{i\pi\omega}{k^2}\frac{d\delta(v_k)}{dv_k} \right] \times \right. \\ & \left(-\frac{v_z}{c} B_\perp \right) \left(-\frac{2v_k}{v_{\text{th},e}^2} + \frac{\epsilon\delta_T}{2v_{\text{th},e}} \left[\cos\theta \left(5 - \frac{2v^2}{v_{\text{th},e}^2} \right) - \frac{2v_k}{v_{\text{th},e}^2} (v_k \cos\theta - v_\perp \sin\theta) \left(7 - \frac{2v^2}{v_{\text{th},e}^2} \right) \right] \right) \\ & \left. + \frac{B_\perp}{ck} \left[-\frac{2v_z}{v_{\text{th},e}^2} - \frac{\epsilon\delta_T}{2v_{\text{th},e}} (v_k \cos\theta - v_\perp \sin\theta) \left(7 - \frac{2v^2}{v_{\text{th},e}^2} \right) \right] \right\} f_{\text{m},e}, \quad (29) \\ f_{\vec{k},i} = & +\frac{ie}{m_i} \left\{ \left[\frac{1}{k}P\frac{1}{v_k} + \frac{\omega}{k^2}P\frac{1}{v_k^2} + \frac{i\pi}{k}\delta(v_k) - \frac{i\pi\omega}{k^2}\frac{d\delta(v_k)}{dv_k} \right] \times \right. \\ & \left(-\frac{v_z}{c} B_\perp \right) \left(-\frac{2v_k}{v_{\text{th},i}^2} + \frac{\epsilon\delta_T}{2v_{\text{th},i}} \left[\cos\theta \left(5 - \frac{2v^2}{v_{\text{th},i}^2} \right) - \frac{2v_k}{v_{\text{th},i}^2} (v_k \cos\theta - v_\perp \sin\theta) \left(7 - \frac{2v^2}{v_{\text{th},i}^2} \right) \right] \right) \end{aligned}$$

$$+ \frac{B_{\perp}}{ck} \left[-\frac{2v_z}{v_{\text{th},i}^2} - \frac{\epsilon\delta_T}{2v_{\text{th},e}} (v_k \cos \theta - v_{\perp} \sin \theta) \left(7 - \frac{2v^2}{v_{\text{th},i}^2} \right) \right] \Big\} f_{\text{m},i} \quad (30)$$

for mode (2).

The Poisson equation is written as

$$ikE_k = 4\pi e \left(\langle f_{\vec{k},i} \rangle - \langle f_{\vec{k},e} \rangle \right). \quad (31)$$

Keeping only the first non-vanishing order in ϵ for mode (1), each term on the right-hand side provides

$$-4\pi e \langle f_{\vec{k},e} \rangle = -\frac{i}{k} \left[\frac{2}{\lambda_D^2} E_k - \epsilon\delta_T \frac{\omega_{\text{pe}}^2}{cv_{\text{th},e}} B_z \sin \theta \right], \quad (32)$$

$$+4\pi e \langle f_{\vec{k},i} \rangle = -\frac{i}{k} \left[\frac{2}{\lambda_D^2} E_k - \epsilon\delta_T \frac{\omega_{\text{pi}}^2}{cv_{\text{th},i}} B_z \sin \theta \right], \quad (33)$$

where ω_{pe} and ω_{pi} are the electron and the ion plasma frequencies, defined by $\omega_{\text{pe}} \equiv \sqrt{(4\pi n_0 e^2/m_e)}$ and $\omega_{\text{pi}} \equiv \sqrt{(4\pi n_0 e^2/m_i)}$, respectively; $\lambda_D \equiv \sqrt{k_B T/(4\pi n_0 e^2)}$ is the Debye length. It is trivial why those equations do not contain $E_{\perp}$ since the transverse component is not constrained from the Poisson equation. Note that the first term in the right hand side for ion is exactly the same as that for electron. Therefore, the contribution of ion to the charge density is non-negligible. On the other hand, the second term for ion is smaller by factor of $\sqrt{m_e/m_i}$ than that for electron. This is the essential point to make possible in-phase acoustic oscillation between electron and ion. Then the spatial charge carried by ion acoustic oscillation is able to be canceled by electron to keep charge neutrality. In the case of pure electron plasma examined by Ramani & Laval (1978), electrons have to keep charge neutrality by themselves. Therefore, the amplitude of acoustic oscillation for electron plasma has to be almost zero. The Poisson equation provides the relation between electric and magnetic fields as

$$E_k = \frac{\epsilon\delta_T}{4} \frac{\sin \theta}{1 + (k\lambda_D)^2} \frac{v_{\text{th},e}}{c} B_z. \quad (34)$$

The term $(k\lambda_D)^2$ in denominator comes from the left-hand side of the Poisson equation and always very small number since the wavelength of the mode which we are interested in is much larger than the Debye length. We, therefore, neglect this term. This corresponds to the so-called Plasma approximation (Chen 1974; Tanaka & Nishikawa 1996) and ensures that the charge neutrality is kept in high accuracy even though the longitudinal oscillation exists. When the wave vector is parallel to the temperature gradient ($\theta = 0$), this equation tells $E_k = 0$ and the mode (1) becomes pure transverse. Compared with the case of pure electron oscillation as in Ramani & Laval (1978), the right-hand side of equation (34) is factor of two smaller, which makes mode (1) more unstable as shown in below. No constraint comes from the Poisson equation for mode (2) since it is a pure transverse mode.

We are now in a position to derive the dispersion relation. The Ampère's law is written as

$$-ikB_z = \frac{4\pi e}{c} \left(\langle v_{\perp} f_{\vec{k},i} \rangle - \langle v_{\perp} f_{\vec{k},e} \rangle \right) + \frac{\partial E_{\perp}}{c\partial t} \quad (35)$$

for mode (1), and

$$ikB_{\perp} = \frac{4\pi e}{c} \left(\langle v_z f_{\vec{k},i} \rangle - \langle v_z f_{\vec{k},e} \rangle \right) + \frac{\partial E_z}{c\partial t} \quad (36)$$

for mode(2). Under the low-frequency condition, the last terms in the right-hand side of equations (35) and (36), those are the displacement current, are negligibly small. We thus neglect these terms. Further, the contribution of ion to the current density is order of $\sqrt{m_e/m_i}$ smaller than that of electron since the current density carried by each particle is proportional to the thermal velocity of each particle. Therefore, the current density carried by ion introduces only a negligible contribution to the dispersion relation. The dispersion relation for the real parts ω_r and the imaginary parts ω_i in the leading order of ϵ is then obtained as follows:

$$\omega_r = \frac{\epsilon \delta_T}{4} k v_{th,e} \cos \theta, \quad (37)$$

$$\omega_i = \frac{\epsilon^2 \delta_T^2}{4\sqrt{\pi}} k v_{th,e} (2 \cos^2 \theta - \sin^2 \theta) - \frac{1}{\sqrt{\pi}} \left(\frac{c}{\omega_{pe}} \right)^2 k^3 v_{th,e} \quad (38)$$

for mode (1), and

$$\omega_r = \frac{\epsilon \delta_T}{4} v_{th,e} k \cos \theta, \quad (39)$$

$$\omega_i = \frac{\epsilon^2 \delta_T^2}{2\sqrt{\pi}} v_{th,e} k \cos^2 \theta - \frac{1}{\sqrt{\pi}} \left(\frac{c}{\omega_{pe}} \right)^2 v_{th,e} k^3, \quad (40)$$

for mode (2).

Since the imaginary part ω_i is of order ϵ^2 , we have checked whether the leading order of the dispersion relation is changed when the second-order distribution functions $f_e^{(2)}$ and $f_i^{(2)}$ are taken into consideration. We have confirmed that these second-order distribution functions only introduce one order higher terms both in real and imaginary parts, and thus main results shown above are not changed.

5. Instability and mode characteristics

The dispersion relations (37), (38), (39) and (40) show that the low-frequency mode, for which the phase velocity of the wave is slower than ion thermal velocity, can exist as long as $\epsilon < \sqrt{m_e/m_i} \sim 0.025$ is satisfied. This is the first confirmation that the Ramani & Laval (1978) type instability can exist beyond the wall of square root of the mass ratio. The dispersion relations are almost the same as those obtained for electron plasma by Ramani & Laval (1978). However, there is a slight difference in imaginary part of the mode (1). The difference comes from the difference in the charge neutrality condition (34). As explained in §4, in our case, the in-phase acoustic oscillation between electron and ion is possible. Hence, the spatial charge carried by ion acoustic oscillation is able to be canceled by electron to keep charge neutrality. However, in the case of pure electron plasma examined by Ramani & Laval (1978), electrons have to keep charge neutrality by themselves. Therefore, the amplitude of acoustic oscillation for electron plasma has to be almost zero.

The characteristics of the instability are summarized as follows. For mode (1), the imaginary part of the wave frequency is positive and the instability sets in, when the direction of the wave vector is within the double cone spanned by $\theta \in [-\theta_{cr}, \theta_{cr}]$ and $\theta \in [\pi - \theta_{cr}, \pi + \theta_{cr}]$, where $\theta_{cr} \equiv \arccos(1/\sqrt{3})$ ($0 \leq \theta_{cr} \leq \pi/2$). For comparison, we re-calculated the dispersion relations for pure electron plasma. In the case of pure electron plasma, a factor of 2 appears in front of the $\sin^2 \theta$ term in the imaginary part (this result is slightly different from Ramani & Laval (1978)'s result.). Therefore, the unstable region in $\vec{k}$ -space in the case of pure electron plasma is somewhat narrower than our case. It shows that acoustic oscillation of ion

is assisting the instability. For mode (2), the unstable mode exists for all direction of the wave vector, although the growth rate decreases as θ increase from 0 to $\pi/2$ for fixed k . Two modes get identical when $\theta = 0$. For both modes, the growth rate is maximum when $\theta = 0$. Since the imaginary part of the wave frequency is the third order polynomial of k with negative coefficient for k^3 , there exists a maximum growth rate $\omega_{i,\max}$. For mode (1), $\omega_{i,\max}$ is given as

$$\omega_{i,\max} \sim \frac{\epsilon^3 \delta_T^3}{12\sqrt{3}\pi} \frac{v_{th,e}}{c} \omega_{pe} |3 \cos^2 \theta - 1|^{3/2}, \quad (41)$$

when

$$k = k_{\max} = \frac{\epsilon \delta_T}{2\sqrt{3}} \frac{\omega_{pe}}{c} |3 \cos^2 \theta - 1|^{1/2}, \quad (42)$$

for $\theta \in [-\theta_{cr}, \theta_{cr}]$ or $\theta \in [\pi - \theta_{cr}, \pi + \theta_{cr}]$. For mode (2),

$$\omega_{i,\max} \sim \frac{\epsilon^3 \delta_T^3}{3\sqrt{6}\pi} \frac{v_{th,e}}{c} \omega_{pe} |\cos^3 \theta| \quad (43)$$

when

$$k = k_{\max} = \frac{\epsilon \delta_T}{\sqrt{6}} \frac{\omega_{pe}}{c} |\cos \theta|, \quad (44)$$

for arbitrary θ . For the wave vector which satisfies the above conditions, the real part ω_r of the wave frequency is obtained as

$$\omega_r \sim \frac{\epsilon^2 \delta_T^2}{8\sqrt{3}} \frac{v_{th,e}}{c} \omega_{pe} |3 \cos^2 \theta - 1|^{1/2} \cos \theta \quad (45)$$

for mode (1), and

$$\omega_r \sim \frac{1}{4\sqrt{6}} \epsilon^2 \delta_T^2 \frac{v_{th,e}}{c} \omega_{pe} \cos^2 \theta \quad (46)$$

for mode (2).

The electric field strength can be related to the magnetic field strength. For mode (1),

$$|E_k| \sim \frac{\epsilon \delta_T}{4} \frac{v_{th,e}}{c} |\vec{B}_k| |\sin \theta|. \quad (47)$$

When $k = k_{\max}$, the electric field perpendicular to the $\vec{k}$ -direction is related to the magnetic field as

$$|E_{\perp}| \sim \frac{\epsilon \delta_T}{4} \frac{v_{th,e}}{c} |\vec{B}_k| |\cos \theta|. \quad (48)$$

For mode (2),

$$|E_z| \sim \frac{\epsilon \delta_T}{4} \frac{v_{th,e}}{c} |\vec{B}_k| |\cos \theta|, \quad (49)$$

when $k = k_{\max}$.

The electric field strength is then order of ϵ smaller than the magnetic field strength. This nature combined with the low-frequency nature of the mode shows that the mode has similarity with the magneto-hydrodynamical mode. However, the magneto-hydrodynamical treatment cannot identify the mode and the instability. For example, the instability is microscopical, in which case the resonance of particles with waves is essential.

The possible reduction factor of the mean free path, in other words the possible lowest value of ϵ , is set by the application limits and assumptions summarized in §2. In the following discussion, we treat the fastest growing mode, that is $\theta = 0$ case. First, the condition (6) provides

$$\epsilon > 4 \left(\frac{1}{L} \frac{c}{\omega_{pe}} \right)^{1/2} \left(\frac{\delta_T}{1.0} \right)^{-3/2}. \quad (50)$$

Second, the age condition (7) provides

$$\epsilon > 2 \left(\frac{H_0}{v_{th,e}} \frac{c}{\omega_{pe}} \right)^{1/3} \left(\frac{\delta_T}{1.0} \right)^{-1}. \quad (51)$$

Third, the collisionless condition is satisfied when $2\pi/k_{\max} < \epsilon \times L$, which yields

$$\epsilon > 4 \left(\frac{1}{L} \frac{c}{\omega_{pe}} \right)^{1/2} \left(\frac{\delta_T}{1.0} \right)^{-1/2}. \quad (52)$$

Finally, the condition that the life time of the interface given by equation (1) must be longer than the age of the cool region, $t_{\text{age}}(\text{cool})$, provides the upper limit on ϵ as

$$\epsilon < 10^{-5} \left(\frac{t_{\text{age}}(\text{cool})}{10^{10}\text{yr}} \right)^{-1} \left(\frac{L}{4\text{kpc}} \right) \left(\frac{k_B T_e}{4\text{keV}} \right)^{-0.5}. \quad (53)$$

6. Application to the cluster of galaxies

6.1. A jam bun model

For the several nearby clusters of galaxies, the existence of at least the two different temperature gas in the central region is spectroscopically confirmed (Ikebe 1995). If the cool gas found in the cluster central region is a single block of the low temperature gas located at the bottom of the cluster gravitational potential well like a jam in a jam bun, the interface between the cool and hot gas can be well described by the pressure equilibrium condition and the results obtained in this paper can be applied. One of the well studied such clusters is the Centaurus cluster (Ikebe et al. 1999). In the following discussion, the cool gas found in the Centaurus cluster is assumed to be well described by the jam bun model. The temperature of the cool region is about $k_B T_1 \sim 1\text{keV}$ and that of the hot region is about $k_B T_2 \sim 4\text{keV}$ in the Centaurus cluster (Ikebe et al. 1999). The factor δ_T is obtained as $\delta_T \sim (T_2 - T_1)/(T_2 + T_1)/2 \sim 1.2$. The size of the region, where the cool component exists, is about the size of the X-ray halo around the elliptical galaxies, say $50\text{--}100h^{-1}\text{kpc}$ (Forman, Junes, & Tucker 1985; Matsushita et al. 1998; Ikebe et al. 1999). Using the best fit values for the electron density profile for the hot phase gas obtained by ASCA results (Ikebe et al. 1999), the electron number density of the hot phase gas at around 100kpc can be estimated as $2.5 \times 10^{-3}\text{cm}^{-3}$. Then the electron mean free path λ_C due to Coulomb collisions is about 2kpc . Hence, the condition (5) implies $L > 2\text{kpc}$, at least when the low-temperature region was formed. The age condition (51) leads to

$$\epsilon > 4 \times 10^{-7} h^{1/3} \left(\frac{k_B T_e}{4\text{keV}} \right)^{-1/6} \left(\frac{n_0}{2.5 \times 10^{-3}\text{cm}^{-3}} \right)^{-1/6} \left(\frac{\delta_T}{1.2} \right)^{-1}. \quad (54)$$

The collisionless condition (52) also provides a lower limit on ϵ as

$$\epsilon > 10^{-7} \left(\frac{n_0}{2.5 \times 10^{-3}\text{cm}^{-3}} \right)^{-1/4} \left(\frac{L}{4\text{kpc}} \right)^{-1/2} \left(\frac{\delta_T}{1.2} \right)^{-1/2}. \quad (55)$$

Therefore, the age condition yields the lower limit on ϵ when $L > 0.25\text{kpc}$ and is compatible with the life time condition (53). On the other hand, the condition (55) provides the lower limit on ϵ when $L < 0.25\text{kpc}$. In this case, for this condition to be compatible with the life time condition (53), the lower limit of the size of the interface region is set as

$$L > 0.2 \left(\frac{t_{\text{age(cool)}}}{10^{10}\text{yr}} \right)^{2/3} \left(\frac{n_0}{2.5 \times 10^{-3}\text{cm}^{-3}} \right)^{-1/6} \left(\frac{k_B T_e}{4\text{keV}} \right)^{1/3} \left(\frac{\delta_T}{1.2} \right)^{-1/3} \text{kpc}. \quad (56)$$

Compared with the cluster pressure scale height, the obtained small values of the lower limit on L both from the age condition and the condition (56) guarantee the application of the pressure equilibrium condition. Therefore, we conclude that the instability found in this paper can be a possible mechanism to inhibit the heat conduction between cool and hot phases in the cluster central region, if the cool gas is a single block of the low temperature gas located at the bottom of the cluster gravitational potential well.

6.2. The raisin bread model

If the cool gas is contained in small blobs and the blobs distribute in the entire cluster central region like raisins in a raisin bread, the obtained results cannot be simply applied. The cooling flow model, which is one of the widely-accepted model to describe the cluster central region (Fabian 1994), assumes that the ICM in the cluster central region is in multi-phase and is composed of many small cool clumps (raisins) which have different temperatures from clump to clump. Even in the two temperature phase model (Ikebe et al. 1999), the raisin bread model is thought to be more plausible to describe the situation in the cluster central region. It is well known that these cool clumps, which is in pressure equilibrium with surrounding hot medium in the cluster central region, oscillates owing to buoyancy force (Balbus 1986; Malagoli, Rosner and Bodo 1987; Balbus and Soker 1989) and that they are fragmented (Nulsen 1986; Balbus 1986) and merge into the surrounding hot medium within a few buoyancy oscillation period, that is $\sim a \text{ few} \times 10^8\text{yr}$ (Hattori and Habe 1990; Malagoli, Rosner and Fryxell 1990; Reale et al. 1991; Yoshida, Hattori and Habe 1991). It has been shown that the magnetic field frozen in the ICM can prevent the buoyancy oscillation of the cool clumps (Nulsen 1986; Loewenstein 1989; Balbus 1991; Hattori, Yoshida and Habe 1995). Therefore, the raisin bread model requires the existence of the magnetic field in the plasma. Nulsen (1986) pointed out that if the Alfvén velocity v_A of the hot medium is larger than the terminal velocity v_t of the cool clumps, that is, if

$$v_A > v_t, \quad (57)$$

the relative motion of cool clumps against the surrounding medium is stopped and the fragmentation of the clumps are prevented. The terminal velocity of a cool clump with a relative density excess of $\delta\rho$ and a size of r is given by $v_t = \sqrt{(\delta\rho/\rho)(r/R)}$, where R is the pressure scale height of the cluster. It was confirmed by the numerical simulation (Hattori, Yoshida and Habe 1995) that this condition works well even for the cool clumps with $\delta\rho/\rho > 1$. It is important to note that even if the magnetic field line is parallel to the gravitational force, the buoyancy oscillation can be prevented by the magnetic tension which is first pointed out by Loewenstein (1989).

It can be shown that the gas pressure equilibrium condition is a good approximation even for the cool clumps in the raisin bread model. Suppose that a cool gas clump is in pressure equilibrium with surrounding medium which is immersed in the background magnetic field. The cool gas clump falls ahead of the surrounding medium owing to its excess weight. As a result, the magnetic field configuration of the surrounding medium is locally disturbed. The infall of the cool clump is stopped, when the force produced

by the disturbed magnetic field acting on the cool clump balance with the buoyancy force. This condition is written as

$$\frac{P_m \delta_B}{r} \sim \delta_\rho \frac{P_g}{R}, \quad (58)$$

where P_m and P_g are magnetic and gas pressures of the hot surrounding medium, respectively; $\delta_B = \delta B/B$ and $\delta_\rho = \delta \rho/\rho$ are deviations of the magnetic field strength and the gas density in the clump, respectively, relative to those of surrounding medium when the force balance is realized. This condition can be rewritten using the plasma β ($\beta \equiv P_g/P_m$) as

$$\delta_B \sim \beta \delta_\rho \frac{r}{R}. \quad (59)$$

From equation (59), the condition (57) can be interpreted as $1 > \delta_B$, and the condition for the possible cool clumps in the fixed- β plasma is obtained as

$$\frac{1}{\beta} > \delta_\rho \frac{r}{R}. \quad (60)$$

In the equilibrium state, the total pressure in the clump must be the same as the total pressure of the surrounding hot medium. This condition provides

$$|\delta P_g| \sim |\delta_B P_m|, \quad (61)$$

in the clump. Therefore, the deviation of the gas pressure relative to the surrounding gas pressure can be estimated as

$$\frac{|\delta P_g|}{P_g} \sim \delta_\rho \frac{r}{R}. \quad (62)$$

From equations (60) and (62), we obtain the constraint on the relative deviation of the gas pressure:

$$\frac{1}{\beta} > \frac{|\delta P_g|}{P_g}. \quad (63)$$

Since the hot gas in the cluster is high- β plasma, say $\beta \sim 10 - 100$ (Kim et al. 1989; Makino 1997), the relative deviation of the gas pressure is at most 10% and very small.

The arguments presented above ensures that the model of the interface adopted in this paper is applicable to the case when the raisin bread model is the good representative for the cluster central region. Therefore, the results obtained in this paper can provide some hint for reduction of the heat conductivity in cooling flow. However, the further analysis including the background magnetic field is required to apply the raisin bread model since the background magnetic field introduces another resonance related to cyclotron frequencies of electron and ion.

7. Discussion

We have shown that there exists low-frequency growing modes driven by a global temperature gradient in electron and ion plasmas. The instability provides a hint for reduction mechanism of the thermal conductivity to explain the observational evidence of co-existence of the multi-phase gas in clusters of galaxies. However, to answer whether the instability really reduces the heat conductivity down to the required level, non-linear saturation level of the instability and the effect of the back ground magnetic field must be examined. Although Levinson & Eichler (1992) has studied the effect of the magnetic field on the Ramani and Laval type instability for pure electron plasma, there is no study which includes ion with back ground magnetic field.

The evolutionary nature of the modes in the interface region can be briefly described as follows. When the low-temperature gas is injected into the intracluster medium, any wave is not excited in the interface region yet. Hence, ϵ is defined by the mean free path due to Coulomb collisions and is order of 1 to 0.1, as shown in §1. At this initial time period, the Ramani & Laval (1978) type instability excites the waves only for electron plasma, since the phase velocity of excited waves is about the electron thermal velocity and thus much faster than the ion thermal velocity. As waves grow, the mean free paths of electron and ion become shorter and ϵ gets smaller. When ϵ goes down to $\sqrt{m_e/m_i}$, ions start to respond the waves, and the modes switch to those described in this paper. Since the growth rate of the instability is also proportional to ϵ^3 and very large at the initial stage, the condition (50) is satisfied. We thus do not have to worry about the disappearance of the interface due to the thermal conduction before the waves grow. The waves must be scattered back at the edge of the interface and must be confined in the interface region, because the waves driven by the instability cannot propagate in the uniform temperature region.

Since the instability creates and grows the magnetic field without any seed field, it provides a new possibility for the origin of the cosmic magnetic field. The Biermann battery effect (Biermann 1950) is the mechanism which has been considered as one of the central mechanism to create the seed magnetic field in our universe (Kulsrud et al. 1997). The Biermann battery effect requires that the direction of the density gradient of the gas is not parallel to the direction of the temperature gradient. However, the instability found in this paper requires only the existence of the temperature gradient. In the astrophysical situation, the situation that two different temperature plasmas are contacting keeping pressure equilibrium can be commonly found compared with the situation that the Biermann battery effect may act. Therefore it may be interesting to examine how the instability found in this paper plays a role on the origin of the cosmic magnetic field.

Finally, it is interesting to note that our results suggest that the realization of the global thermal equilibrium is postponed by the local instability which is induced for quicker realization of local thermal equilibrium state in plasmas. This result is of course not trivial. Once the global thermal equilibrium is achieved, the local thermal equilibrium is realized simultaneously. Therefore, letting electrons be free to conduct thermal energy without any instability is the fastest way to achieve thermal equilibrium. However, the plasma prefers restoring the local deviation of the velocity distribution function from the Maxwell-Boltzmann and excites waves by the instability. As a result, the realization of the local thermal equilibrium is also postponed. The plasma be hurried into an error.

Authors would like to thank Masahiro Hoshino, Rudolf Treumann, Yousuke Itoh, Yutaka Fujita, Masahide Iijima for many instructive and fruitful discussion on this topics.

REFERENCES

- Balbus, S.A. 1986, *ApJ*, 303, L79
- Balbus, S.A., & Soker, N. 1989, *ApJ*, 341, 611
- Balbus, S.A. 1991, *ApJ*, 372, 25
- Biermann, V. L. 1950, *Z. Naturforsch.*, 5a, 65
- Chapman, S., & Cowling, T. G. 1960, *The Mathematical theory of non-uniform gases*, (Cambridge University Press, Cambridge)
- Chen, F. F. 1974, *Introduction to Plasma Physics*, (Plenum Press, New York)

- Fabian, A. C. 1994, *Ann.Rev.Astron.Astrphys.*, 32, 277
- Forman, W., Jones, C., & Tucker, W. 1985, *ApJ*, 293, 102
- Fried, B. D. 1959, *Phys. Fluid.*, 2, 337
- Fukazawa, Y., Ohashi, T., Fabian, A. C., Canizares, C.R., Ikebe, Y., Makishima, K., Mushotzky, R. F., & Yamashita, K. 1994 *PASJL* 46, L55
- Hattori, M., & Habe, A. 1990, *MNRAS*, 242, 399
- Hattori, M., Yoshida, T., & Habe, A. 1995, *MNRAS*, 275, 1195
- Hattori, M., Kneib, J.-P., & Makino, N. 1999, *PTP Suppl.*, 133, 1
- Ikebe, Y. 1995, PhD. Thesis, (University of Tokyo)
- Ikebe, Y., Makishima, K., Fukazawa, Y., Tamura, T., Xu, H., Ohashi, T., & Matsushita, K. 1999, preprint
- Kim, K.T., Kronberg, P.P., Giovannini, G., & Venturi, T. 1989, *Nat*, 341, 720
- Kulsrud, R. M., Cen, R., Ostriker, J. P., & Ryu, D. 1997, *ApJ*, 480, 481
- Levinson, A., & Eichler, D. 1992, *ApJ*, 387, 212
- Loewenstein, M. 1990, *ApJ*, 349, 471
- Makino, N. 1997, *ApJ*, 490, 642
- Makishima, K. 1997, in *Imaging and Spectroscopy of Cosmic Hot Plasmas*, eds. F. Makino & K. Mitsuda (University Academy Press, Tokyo), p137
- Malagoli, A., Rosner, R., & Bodo, G. 1987, *ApJ*, 319, 632
- Malagoli, A., Rosner, R., & Fryxell, B. 1990, *MNRAS*, 247, 367
- Matsushita, K., Makishima, K., Ikebe, Y., Rokutanda, E., Yamasaki, Y. N., & Ohashi, T. 1998, *ApJ*, 499, L13
- Melrose, D. B. 1986, *Instabilities in space and laboratory plasmas*, (Cambridge University Press, Cambridge)
- Nulsen, P. J. E. 1986, *MNRAS*, 221, 377
- Pistner, S., Levinson, A., & Eichler, D. 1996, *ApJ*, 467, 162
- Ramani, A., & Laval, G. 1978, *Phys. Fluid.*, 21(6), 980
- Reale, F., Rosner, R., Malagoli, A., Peres, G., Serio, S. 1991, *MNRAS*, 251, 379
- Sarazin, C.L. 1988, *X-ray emission from clusters of galaxies*, (Cambridge University Press, Cambridge)
- Soker, N., & Sarazin, C.L. 1990, *ApJ*, 348, 73
- Tanaka, Y., Inoue, H., & Holt, S. S. 1994, *Publ.Astron.Soc.Jpn.*, 46, L37
- Tanaka, M., & Nishikawa, K. 1996, *Physics of high temperature plasmas*, (Maruzen, Tokyo)
- Weibel, E. S. 1959, *Phys.Rev.Lett.*, 2, 83
- White, R. A., & Sarazin, C. L. 1987, *ApJ*, 318, 612
- Yoshida, T., Hattori, M., Habe, A. 1991, *MNRAS*, 248, 630

Fig. 1.— A sketch of one-dimensional distribution of temperature $T(x)$ in the interface region. T_1 and T_2 are the temperatures in the cool and hot regions, respectively. Also shown are the number density distribution $n(x)$ (dashed line) and the distribution of pressure (dotted line), which is constant along the x -direction.

